



Linton, E., Walkden, A., Steeples, L. R., Bhargava, A., Williams, C. E. M., Bailey, C., Quhill, F., & Kelly, S. (2018). Retinal burns from laser pointers: a risk in children with behavioural problems. *Eye*, 33, 492-504. <https://doi.org/10.1038/s41433-018-0276-z>

Peer reviewed version

License (if available):  
CC BY-NC

Link to published version (if available):  
[10.1038/s41433-018-0276-z](https://doi.org/10.1038/s41433-018-0276-z)

[Link to publication record in Explore Bristol Research](#)  
PDF-document

This is the accepted author manuscript (AAM). The final published version (version of record) is available online via Nature at <https://doi.org/10.1038/s41433-018-0276-z>. Please refer to any applicable terms of use of the publisher.

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:  
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

# **Retinal burns from laser pointers, a risk in children with behavioural problems.**

Keywords: CHILDREN, EYE INJURY, LASER, RETINA, SELF-INJURIOUS BEHAVIOUR

**Linton E<sup>1</sup>, Walkden A<sup>2</sup>, Steeples LR<sup>3</sup>, Bhargava A<sup>2</sup>, Williams C<sup>3</sup>, Bailey C<sup>3</sup>, Quhill FM<sup>4</sup>, Kelly SP<sup>1</sup>**

<sup>1</sup> Department of Ophthalmology, Bolton Hospitals NHS Foundation Trust

<sup>2</sup> Department of Ophthalmology, Lancashire Teaching Hospitals NHS Foundation Hospital

<sup>3</sup> Bristol Eye Hospital, University Hospitals Bristol NHS Foundation Trust

<sup>4</sup> Department of Ophthalmology, Sheffield Teaching Hospitals NHS Foundation Trust

*Correspondence to:* Mr Simon P Kelly, FRCOphth,

Eye Dept, Bolton Hospitals NHS Foundation Trust, Bolton. BL4 0JR. England

**Acknowledgment:** We thank Dr Colin Swift, Medical Physics and Engineering Dept, The Christie NHS Foundation Trust, Manchester for testing the laser pointer and comments on the manuscript. We thank Dr John O'Hagan, Laser and Optical Radiation Dosimetry Group, Public Health England for comments on the manuscript.

**Conflicts of interest:** The authors declare no conflicts of interest of relevance.

**Research funding:** No special funding

# Retinal burns from laser pointers

## ABSTRACT

**Objective:** To explore self-inflicted retinal burns from laser pointers in children.

**Methods:** Literature review of laser pointer retinal injuries in childhood and online survey of UK Consultant Ophthalmologists. A cohort of local children with self-inflicted injury are described. The matter is topical. We review progress in recent legislation and policy change in the UK

**Results:** Four of 77 case reports of laser burns in childhood analysed reported psychological or behavioural issues. Three of 4 children in our cohort had such issues. Delay in diagnosis occurred in two of our patients. Structural retinal damage persisted for over 12 months in all 4 children (7 eyes). Our survey of UK ophthalmologists found 159 cases of injury (85% male), 80% under 20 years of age. The majority of the laser pointers were purchased online. Many patients (36%) suffered moderate vision loss (6/18 to 6/60 Snellen), while 17% (at least 11 patients) suffered severe vision loss (<6/60 Snellen).

**Conclusion:** We highlight the risk of macular damage and vision loss from handheld lasers specifically in children with behavioural, learning or mental health issues. The diagnosis may be difficult or delayed in such children. In children with uncertain macular changes, ophthalmologists should explore the history for possible instances of exposure to handheld lasers pointers. Regulatory authorities and manufacturers of handheld lasers need to be aware of the risk to children. Furthermore, there is a need to better inform parents, carers and teachers of the risk of ocular self-injury from such lasers pointers

# Retinal burns from laser pointers

## 1 INTRODUCTION

Laser pointers (sometimes termed laser pens) are handheld laser devices intended for pointing out objects or locations, including for demonstration and amusement purposes. Such lasers should have minimal risk of causing harm to vision. However, retinal injury from laser pointers is causing concern due to the wider availability of more powerful and cheaper laser pointers. The authors have encountered both adults and children with such injuries, including self-inflicted retinal injury from the misuse of high-powered handheld laser pointers. To explore this further in childhood we undertook both qualitative and quantitative ('mixed methods') research and also met with stakeholders in the UK. The material presented herein includes a literature review, a survey of UK ophthalmologists, clinical follow up of 7 eyes of 4 local children with self-inflicted laser burns diagnosed in UK hospital practice and an update of our engagement with stakeholders and policy makers.

## ***METHODS***

### **1 Literature review.**

We located all reports of laser pointer injury available on MEDLINE (on Ovid from 1966) and EMBASE (on Ovid from 1980) and ISI Web of Science (from 1990).

Keywords and MESH terms for 'laser pointer' and 'retina' or its similes were used. The final list of titles and abstracts was screened by two reviewers (EL and AW) and full publications were obtained where articles were thought to be potentially relevant.

Bibliographies of included studies and review papers were screened to identify other relevant studies. The literature search is accurate and up to date as of 19<sup>th</sup> March 2018. We searched for reports of self-inflicted laser burns where children were involved and then systematically explored for any psychological and behavioral features

## **Retinal burns from laser pointers**

recorded in such reported childhood cases. We excluded patients described as being over 18 years old at time of injury and studies where the full articles were not available in English.

### **2 Survey of UK Ophthalmologists.**

An online survey of UK consultant ophthalmologists was undertaken in January 2016 by one of our senior authors to explore their experience of laser pointer injury. A brief online survey was emailed to 990 consultant ophthalmologists in the UK, asking whether they had encountered a patient who suffered macular injury due to misuse of a handheld laser device.

Ophthalmologists who gave a positive answer were also asked: the number of such laser pointer burn patients they had encountered; ages and gender of patients; whether the injury was accidental, self-inflicted or deliberate; the power and colour of the laser beam and where purchased; visual outcome and optical coherence topography (OCT) and visual field evidence. To keep the survey brief and encourage completion, ophthalmologists who indicated seeing more than 2 patients were only asked to provide the details of the most and least affected patients. The data was analysed based on fully completed surveys.

### **3 Case Series.**

A convenience sample of four children (7 eyes) with self-inflicted retinal injury from laser pointers who presented to hospitals in Bolton, Bristol and Preston within a 12 month period and who have over 12 months follow up are presented. Informed parental consent for publication of clinical details and images was obtained from all children in this cohort.

### RESULTS

#### Literature review

In the literature review we located 84 cases of handheld laser burns in children age 18 years or younger reported on 19/03/2018.<sup>1-46</sup> (**Table1:Supplementary Material** ).This includes a case series that the senior authors (SPK and FMQ) previously provided.<sup>12</sup> Within these reports we systematically located one child with a pre-existing diagnosis of attention deficit hyperactivity disorder (ADHD), a second child had known learning difficulties and the third who was undergoing psychological treatment following a road traffic accident.<sup>8-9,17</sup> In one further case report we detected that a young person was referred for psychiatric evaluation following retinal injury from self-harming behaviour with handheld lasers.<sup>38</sup> We acknowledge a case report of laser maculopathy in a twenty year old man in France with schizoid personality, but this is excluded from Table 1 which highlights cases of children.<sup>21</sup> Two further abstracts were identified of laser eye injuries in children but the full articles were not available in English, therefore limited information is included in Table 1 and we cannot exclude any contributing psychological or behavioural problems in these children.<sup>22,29</sup>

#### Survey of UK Ophthalmologists

The survey submitted to 990 Consultant Ophthalmologists in the UK, using a 'mailing' database of email addresses of UK NHS Consultants, by one of the senior authors (FMQ) had a response rate of 15.5% and identified 159 cases of macular injury. Many injuries occurred within the year preceding the survey (54%) with most of the affected patients (80%) under 20 years of age or male (85%).

## Retinal burns from laser pointers

Most laser pointers were reported as having been purchased online. Many patients (36%) suffered moderate vision loss (6/18 to 6/60 Snellen), while 17% suffered severe vision loss (<6/60 Snellen). Visual acuity was not affected in 15% of cases.

Many of the injuries happened due to lack of awareness of the danger, and were either self-inflicted (35%) or caused by a third party (36%). There were no cases of assault reported. No relevant results on the colour of laser beam was provided. The power of known devices exceeded 50 mW in 33% of cases. The survey has been presented as a poster.<sup>47</sup>

**Case series.** We describe 4 local children (7 eyes) with self-inflicted retinal damage from handheld laser pointers with more than 12 months follow up. All showed persistent outer retinal lamellar layer defects on spectral domain ocular coherence tomography (SD-OCT). Three children had a history of mental health or psychological challenges. All cases presented to our 3 hospitals within a 12 month period.

### **Case 1**

An 11 year old male, with a diagnosis of pathological demand avoidance (PDA) and migraine, presented to a community optometrist with a two day history of a black spot in the central vision of his right eye. The optometrist reported that the best corrected visual acuity (BCVA) in the right eye was reduced to 6/10 Snellen having been normal at a prior visit. Left was 6/5 Snellen. New pigmentary changes at both macula were observed by the optometrist and referral was made to the hospital eye service (HES). He was taking pizotifen prescribed for migraine. There was no relevant past ocular, medical or family history. Six weeks later, in the hospital eye service (HES) review, he described a persistent 'blur' in the central vision of the right eye. Unaided VAs were 6/9 right and 6/5

## Retinal burns from laser pointers

left. Pigmentary changes were noted at the central macula in both eyes. Spectral domain optical coherence tomography (SD-OCT) and imaging revealed bilateral outer lamellar layer defects. (**Figure 1**). Electrodiagnostic tests were normal but with limited co-operation. His mother accompanied him for all HES visits. In due course and following direct questioning he admitted to constructing a device made from Lego™ consisting of a laser pointer with a condensing lens used just prior to the onset of visual symptoms. The patient's mother revealed that she had purchased the laser pointer online for him. During follow up he reported symptomatic improvement and the VA remained stable. Centre involving structural defects at both macula persisted on clinical examination and OCT imaging to most recent follow up 24 months later. (**SUPPLEMENTARY IMAGE 1**). The mother confirmed the laser had been purchased from a well-known UK online retailer and was still available for online purchase a year following the incident.

### Case 2

A 13 year old male with attention deficit disorder (ADD) presented to the Emergency Medicine Department accompanied by his mother complaining of visual disturbance after staring into the beam from a toy laser for a few hours earlier that day. The patient stated that the toy laser belonged to a friend but the injuries were self-inflicted. The BCVA was 6/60 in the right eye improving to 6/36 with pinhole, and 6/12 in the left eye. SD-OCT images on presentation showed full-thickness hyper-reflective damage involving both fovea (**Figure 2**). The patient was on methylphenidate 57mg daily treatment for ADD and was known to Child and Adolescent Mental Health Services (CAMHS). He attended mainstream school with additional classroom support but was not classified as having special educational needs, with no statement of educational needs undertaken previously. Six weeks later, his BCVA had improved to 6/12 right and 6/9 left. An improvement in SD-OCT images was observed, notably an improvement in inner retinal layers. However, the centre



## Retinal burns from laser pointers

124 involving outer lamellar layer defects on OCT and fundus changes persisted throughout 24  
125 months of follow up but decreased.

### 126 **Case 3**

127 A 15 year old female with no past medical or psychological history was referred following a  
128 routine sight test where new discrete pigmentary changes at the right fovea were  
129 observed. The patient was asymptomatic. Her past ocular history, medical and family  
130 history were unremarkable. The unaided VA was 6/7.5 in both eyes. Two full thickness  
131 centre involving round scars at the right fovea were observed and a third slightly eccentric.  
132 SD-OCT revealed defects in the ellipsoid zone in the outer retina in these lesions (**Figure**  
133 **3**). The patient admitted to being involved in a 'competition game' with three other children  
134 about two years previously in the home. The 'game' consisted of ascertaining which child  
135 could withstand a green laser beam in one eye for the longest time. She recalled shining  
136 the laser into her right eye for short duration, perhaps 10 seconds twice. The laser pointer  
137 had been purchased online by the patient's mother.

138 The laser pointer responsible was retrieved from the family and sent for analysis. The  
139 analysis found the laser pointer was of wavelength 532nm with an average power of  
140 47mW, making it a Class 3B laser. The label on the laser pointer incorrectly stated that it  
141 was "Class II" with a maximum output less than 1mW. (**SUPPLEMENTARY IMAGE 2**). At  
142 latest follow up, at 24 months the macular changes persisted with 6/6 Snellen in each eye.  
143 **SUPPLEMENTARY IMAGE 3**).

### 144 **Case 4**

145 A 12 year old boy was referred with a several month history of reduced vision in both eyes.  
146 He had a history of expressive and receptive language impairment and was attending a  
147 specialist school for children with cognitive impairment and disturbed behaviour. He was

## Retinal burns from laser pointers

under CAMHS for anger and behavioural problems. There was no relevant past ocular or family history. The presenting BCVA were 6/30 right eye and 6/75 left eye. Colour vision was reduced, with only 4/17 Ishihara plates correctly identified in the right eye and 9/17 in the left eye. Bilateral multifocal macular pigmentary changes were noted. **(Figure 4)**. To investigate abnormal visual function, electrodiagnostic tests and magnetic resonance imaging (MRI) of the brain and orbits were performed, both of which were normal.

At subsequent follow up and on direct questioning, the boy revealed he had been playing with laser pointers at school, particularly in games involving looking directly into the beam of the laser pointer. His BCVA at 12 months follow-up was 6/19 right eye and 6/48 left eye. Fundus examination showed irregular pigmentation at the right macular and a scar at the left macular. OCT scan showed small, round, punched-out lesions more frequent in the left than the right macula.

## DISCUSSION

Retinal burns from handheld laser pointers are an important and increasingly topical public health issue. Such devices are becoming more powerful, less costly, are often incorrectly labeled, and can be easily purchased online. Furthermore, there is increasing apprehension for aviation safety following suspected retinal injuries to commercial airline pilots falling victim to laser attacks.<sup>19,27</sup> Lee et al reported that young males were the most frequent group reported to sustain handheld laser pointer injuries in reports from 1999-2014.<sup>2</sup> Our survey of UK ophthalmologists supports these findings, with results showing 85% of reported cases were male and 80% of all patients were under 20 years of age.<sup>47</sup> Our literature review also concurs with these findings, with 73% of cases being young males. While laser burns, including self-inflicted, can affect adults it is opined that children

## **Retinal burns from laser pointers**

are at greater risk of laser pointer injuries than adults as they are intrigued by their appearance, and lack protective mechanisms of blinking and gaze aversion that adults exhibit and furthermore have clear ocular media which provides little protection from laser injury.<sup>2,6-7</sup> The majority of the laser pointer injury cases encountered by the UK ophthalmologists were reported as having occurred from laser pointers that had been purchased online. Clinical management of laser-induced retinal injuries is anecdotal, on occasion oral corticosteroids have been prescribed.<sup>2,7,13</sup>

In our survey of UK ophthalmologists, the reported visual acuity in affected patients was reported as 6/18-6/60 in 36% and worse than 6/60 in 17% of cases.<sup>47</sup> In our literature review the visual acuity at presentation was 6/18-6/60 in 36% and worse than 6/60 in 28% of cases. The final visual acuity, where reported, was 6/18-6/60 Snellen acuity in 24% and worse than 6/60 in 5% of cases.

We acknowledge a recent review by Birtel et al which identified 111 patients of unstated ages with laser pointer eye injuries in the literature.<sup>5</sup> They found highly variable retinal injuries across the literature, including; macular holes, retinal haemorrhage and on OCT imaging disruption of retinal pigment epithelium, outer retinal hyper reflectivity and disruption of outer retinal layers. That review did not document patient factors or patient age or if the injury was self-inflicted.

## **Classifications and misclassification of lasers.**

The revised UK classification of laser products consists of eight categories; Class 1, 1C, 1M, 2, 2M, 3R, 3B and 4, with Class 4 lasers being the highest radiation hazard.<sup>48</sup> The World Health Organisation (WHO) stated in 1998 that “laser pointers higher than class 2 are considered too powerful for general use as laser pointers and present an unacceptable

## **Retinal burns from laser pointers**

risk in the hands of consumers because they may cause eye injury”.<sup>49</sup> Class 2 laser products have a maximum power of 1 milliwatt and fall within the visible wavelength range 400-700nm. In 2014 Public Health England advised “the sale of laser products to the general public for use as laser pointers should be restricted to Class 1 or Class 2 devices” and further advised “toys should be class 1 or of such low output that they do not need to be classified”.<sup>50</sup> In the United States (US) the Food and Drug Administration (FDA) are: class I, IIa, II, IIIa, IIIb and IV with increasing numbers corresponding to higher output power.<sup>51</sup> FDA permits laser pointers with a maximum power of 5 milliwatts (class IIIa) in the visible wavelength region of approximately 400-710 nanometers.<sup>52</sup> However, handheld laser pointers are widely available to purchase online, often do not conform to such regulations or carry appropriate labelling of the laser power or carry warnings with regard to the ocular risk involved. There are reports of these devices being misclassified and found to have a higher output than stated when objectively tested.<sup>2,3,9,11</sup> Incorrect labelling increases ocular hazards; a consumer or parent may think that a Class 2 laser will be safe – but if in reality the device is a Class 3B then the risk will be far greater than anticipated. Recent publications have highlight concerns of incorrect labelling of lasers in the USA, Australia and UK.<sup>53-55</sup> Case 3 in our series is a further example of misclassification. The parents of the children in our series reported that they were unaware of the ocular risks of children misusing laser pointers. This also chimes with other case reports. Lastly, in some cases the parents were unaware that their child was in possession of such devices.<sup>2</sup>

## **Classification and misclassification of laser retinal injury in children**

Diagnosis of laser pointer retinal injuries in childhood can be difficult, as children and parents may be hesitant to admit to use and purchase of such devices. Additionally, laser retinal injuries may have similarities in clinical appearance to other retinal disorders and

## Retinal burns from laser pointers

lead to misdiagnosis, delayed diagnosis and unnecessary investigations or treatment. A recent case series from Moorfields Eye Hospital reported that 5 of the 16 children with laser injury were initially suspected to have macular dystrophies which delayed their diagnosis.<sup>31</sup> Cases 1 and 4 in our series were also initially similarly mistaken as such. We are aware of another case locally being mistaken for macular inflammation. However, the changes seen on SD-OCT imaging namely focal disruption of the ellipsoid zone are diagnostic of photic maculopathy.<sup>2-4,7,9,11,13,39</sup> The recognition of such outer retinal layer defects should prompt a thorough history to enquire if the child has been exposed to a beam from a laser pointer or sun gazing. Zhang et al also commented on similarities between photic macular injuries and macular genetic conditions and opined laser pointer burns patients may improve over time whereas genetic conditions do not.<sup>15</sup> We noted some improvement in the 7 eyes studied but all had centre involving structural damage on SD-OCT persisting after a year or more of follow up. It has been opined that it may be possible to differentiate between self-inflicted and third-party induced laser retinal injuries on SD-OCT imaging. Bhavsar et al reported that self-inflicted laser injuries had a streak-like appearance, whereas injuries caused by others tended to be discrete lesions in close proximity to the fovea.<sup>4</sup> Our study does not confirm this impression as we saw discrete injury in the presence of self-inflicted injury. A recent report of 4 children suggested that the most significant variables predictive of retinal injury in laser pointers are the amount of energy delivered by the laser, duration of exposure and location of retinal involvement.<sup>44</sup> The Moorfields study of children added a proposed classification of severity of laser burn structural damage which we welcome.<sup>31</sup>

## Retinal burns from laser pointers

### **Behavioural and psychological issues in children with self-inflicted injury**

Neither of two recent case series of childhood laser retinal burns or a recent literature review of cases of any age reported any children's co-existent behavioural profiles or whether self-injurious behaviour was a factor.<sup>31,44,5</sup> Similarly in the literature review we undertook searching systematically for such matters in children few reports gave details of children's general or psychological status. We opine that many authors were either unaware of children's behavioural issues or else did not report such details, including absence of any behavioral matters, in their case reports. In any event although we found a small number of case reports that noted existing psychological, psychiatric, behavioural or learning problems in those affected, to our knowledge no reports to date highlight the risk of handheld laser possession in such children or explore a relationship between these diagnoses and laser eye injuries.<sup>8,9,17,38</sup> Case 1 in our series had a diagnosis of pathological demand avoidance (PDA) syndrome. Newson et al described children with PDA as having a resistance and avoidance of demands as well as impulsive and obsessive behaviour and suggested it be a clinical entity in its own right rather than a sub-type of autism.<sup>56</sup> The first systematic comparison of PDA and autism spectrum disorders in 2014 reported that children with PDA showed characteristics of both autism - such as peer problems - as well as traits of conduct disorders such as anti-social behaviour.<sup>57</sup> In our Case 2, the patient had a diagnosis of attention deficit disorder (ADD) , also known as attention deficit hyperactivity disorder (ADHD). Children with ADD/ADHD exhibit behavioural problems and inattention, hyperactivity or impulsivity.<sup>58</sup> In our opinion the common themes of impulsive, obsessive behaviours and a resistance to following instructions puts children with such conditions at risk of self-injurious behaviour and importantly more so if they are in possession or playing unsupervised with objects such as powerful handheld lasers. Our third child did not have any diagnosed mental health problems but did take part in a 'game' that exposed her to direct laser pointer exposure for

## **Retinal burns from laser pointers**

whatever reason. Case 4 had complex behavioural challenges. Two of our 4 children were linked to CAMHS services.

Self-injurious behaviour (SIB) is considered “a class of behaviours, which the individual inflicts upon his/herself that have the potential to result in physical injury”.<sup>59</sup> Weiss explains the subtle comparison between individuals with developmental disabilities unknowingly behaving in a way that leads to harm, and those who set out with an intent to hurt themselves, for example in attempts to take their own life.<sup>60</sup> Self-injurious behaviour has an estimated prevalence of 35-60% amongst people with autism.<sup>61</sup> Self-injurious trauma to the eye is recognised in children with autism and related conditions. Patton reviewed the relevant literature in 2004 and reported that ‘head-banging’ was a common mechanism of ocular injury in children with autism.<sup>62</sup> Very recently, Lee et al reported three case of bilateral cataract following self-inflicted trauma in children with autistic spectrum disorder.<sup>63</sup> Our report highlights another novel ocular self-injurious behaviour in such individuals.

## **Regulation of laser pointers**

Recent editorials by Marshall et al and Bartsch et al provided perspectives on the regulation and safety and hazards of laser pointers from a UK and US viewpoint.<sup>26,64</sup> The review by the Swedish Radiation Safety Authority of 46 cases from the world literature of laser pointer burns is useful as severity and mechanism of injury where known are outlined in that report.<sup>65</sup> In our clinical experience powerful handheld laser pointers in the hands of children with behavioural, learning and or mental health problems is a dangerous risk. We thus wish to draw attention to this hazard. The matter is relevant for parents and regulators. Importantly the classification of laser pointers in various jurisdictions and the advice by Public Health England in the UK does not take into account the potential for ocular harm from prolonged self-inflicted exposure, as occurred in the children reported herein. With

## **Retinal burns from laser pointers**

298 regard to retinal hazards, labels seem designed for laboratory scientists and not  
299 necessarily for the general public and importantly labeling may not reflect the true class of  
300 the laser –as misclassified. A word such as “Class 3R” means little to the non-expert. The  
301 public may falsely assume that these ‘toys’ are safe as they are approved for general sale.  
302 Self-inflicted injury at close range in children and from misclassified laser pointers adds to  
303 our concern as does the increasing availability of cheaper and more powerful handheld  
304 lasers.

305 The senior authors (SPK and FMQ) have alerted the Royal College of Ophthalmologists  
306 and the Royal Society for the Prevention of Accidents on our concerns and attended a  
307 workshop on the matter hosted by Public Health England (PHE). The UK government  
308 reviewed the evidence of harm to children and risk to pilots following a multi-agency  
309 meeting in February 2016 before deciding it’s approach to tackling this mounting ocular  
310 public health problem. One of the senior authors (FMQ) used the data from the recent  
311 online survey of UK ophthalmologists to inform that multi-agency meeting.<sup>47</sup> Following  
312 this multi-agency meeting PHE launched an online health awareness video following the  
313 concern surrounding ocular hazards from laser pointers.<sup>66</sup> We welcome that video and  
314 publicity about laser pointers by some local trading standards authorities.<sup>67</sup> In May 2018,  
315 the Laser Misuse (Vehicles) Act gained Royal Assent. Under this new legislation  
316 individuals who target drivers of trains, buses, boats or planes can be jailed for up to five  
317 years, and the previous cap on the maximum fine of £2500 has been lifted.<sup>68</sup> The  
318 Government Department for Business, Energy and Industrial Strategy ran a Call for  
319 Evidence on Laser Pointers in 2017 and published their response in January 2018.<sup>69</sup> This  
320 document summarises the four steps the Government will take in reaction to the call for  
321 evidence; provide additional support for enforcement activities around the import of high  
322 powered lasers, encourage more effective voluntary labelling of laser pointers, promote  
323 public awareness on the hazards of laser pointers particularly eyesight and address pilots



## Retinal burns from laser pointers

concerns via the aforementioned Laser Misuse (Vehicles) Act 2018.<sup>68,69</sup> We believe that this mixed methods publication assists promoting awareness of a specific ocular public health concern in children in addition to the known wider concerns including for adults. Furthermore the conviction and sentencing in 2016 of an individual for the sale of a laser pointer that caused eye injury in a child was an important step by UK authorities in the enforcement of the regulations surrounding sale of laser products.<sup>70</sup> However we remain concerned about online sale of powerful laser pointers.

The strengths of this mixed methods contribution include its addition to the public health debate and literature by highlighting the risks of retinal burns from laser pointers in children - particularly with respect to children with behavioural problems - and our engagement with UK laser safety stakeholders. We assessed the number, age and gender and visual outcomes of patients with laser injury encountered via UK consultant ophthalmic colleagues using an online survey. A limitation was the poor response rate and thus data so obtained does not provide the true incidence and clinical features of such cases; this which would require formal case finding such as the British Ophthalmic Surveillance Unit (BOSU) undertakes. Our case series is small but has over 12 months follow up data. We are of the opinion that further formal public health case finding and surveillance research is warranted to assess the epidemiology of retinal laser pointer burns and the profile and outcomes of patients who sustain such injury. Cohort studies from hospital eye clinics would be of merit to provide information on OCT biomarkers and prognosis. Such matters may be complicated by the issues that parents may not be aware of their children having laser pointers and or families may be reluctant to disclose such information even where known.

## Retinal burns from laser pointers

348

349

### 350 **IMPLICATIONS FOR POLICY**

351

352 The recognition by UK Government for the need for more robust regulation of the  
353 importation and sale of laser pointers, including online sales is reassuring as is the Laser  
354 Misuse (Vehicles) Act 2018 which was recently given Royal Assent.<sup>67</sup> There is a need for  
355 ophthalmologists to closely question all patients especially children with retinal outer  
356 lamellar layer defects (best appreciated on SD-OCT imaging) for any history of laser  
357 pointer exposure before considering further tests for macular disorders. Importantly there  
358 is a need for increased public awareness and education of the ocular hazards of laser  
359 pointers.<sup>69</sup> In particular, parents, and especially parents of children with conditions that  
360 may increase risk of self-injurious behaviour should be aware powerful and often  
361 incorrectly classified handheld lasers pointers can be dangerous to sight. Specifically the  
362 availability of high powered and also mislabeled laser pointers remains a concern.  
363 Because such lasers are readily available, children likely to self-harm may be at greater  
364 risk of shining laser beam into their eyes, perhaps for longer periods of time. We urge the  
365 manufactures of handheld laser pointers and their vendors to consider our concerns. We  
366 urge the regulators, manufacturers and distributors of laser pointers -including online  
367 merchants- to be more vigilant given this novel concern of vision loss in at risk children.

## Retinal burns from laser pointers

### REFERENCES

1. Yui G, Itty S, Toth CA. Ocular Safety of Recreational Lasers. *JAMA Ophthalmology* 2014; 132:245-246.
2. Lee GD, Bauman CR, Lally D, Pitcher JD, Vander J, Duker JS. Retinal Injury after inadvertent handheld laser exposure. *Retina, The Journal of Retinal and Vitreous Diseases* 2014; 34:2388-2396.
3. Alsulaiman SM, Alrushood AA, Almasaud J, Alzaaidi S, Alzahrani Y, Arevalo JF, Ghazi NG, Abboud EB, Nowilaty SR, Al-Amry M, Al-Rashaed S. High-power handheld blue laser-induced maculopathy: The results of the King Khaled Eye Specialist Hospital Collaborative Retina Study Group. *Ophthalmology* 2014; 121:566-572.
4. Bhavsar KV, Wilson D, Margolis R, Judson P, Barbazetto I, Bailey FK, Cunningham ET. Multimodal imaging in handheld laser-induced maculopathy. *American Journal of Ophthalmology* 2015; 159:227-231.
5. Birtel JM, Harmening WU, Krohne TG, Holz F, Charbel Issa P, Herrmann P. Retinal Injury Following Laser Pointer Exposure: A Systematic Review and Case Series. *Deutsches Arzteblatt International* 2017; 114(49):831-837.
6. Dhoot DS, Xu D, Srivastava S. High-powered laser pointer injury resulting in macular hole formation. *Journal of Paediatrics* 2014; 164:668.
7. Dirani A, Chelala E, Fadlallah A, Antonios E, Cherfan G. Bilateral macular injury from a green laser pointer. *Clinical Ophthalmology* 2013; 7:2127-2130.
8. Fujinami K, Yokoi T, Hiraoka M, Nishina S, Azuma N. Choroidal neovascularisation in a child following laser pointer-induced macular injury. *Japanese Journal of Ophthalmology* 2010; 54:631-633.

## Retinal burns from laser pointers

9. Hanson JVM, Sromicki J, Mangold M, Golling M, Gerth-Kahlert, C. Maculopathy following exposure to visible and infrared radiation from a laser pointer: a clinical case study. *Documenta Ophthalmologica* 2016; 132:147-155.
10. Mainster MA, Stuck BE, Brown J. Assessment of alleged retinal laser injuries. *Archives Ophthalmology* 2004; 123:1210-1217.
11. Noble C, Blice J. Permanent retinal injury from recreational laser pointer. *Military Medicine* 2015;180:378-380.
12. Raoof N, Chan TKJ, Rogers NK, Abdullah W, Haq I, Kelly SP, Quhill FM. 'Toy' laser macular burns in children. *Eye* (2014); 28:231-234.
13. Turaka K, Bryan JS, Gordon AJ, Reddy R, Kwong HM Jr., Sell CH. Laser pointer induced macular damage: case report and mini review. *International Ophthalmology* 2012; 32:293-297.
14. Xu M, Lin T. Bilateral central scotomas following laser pointer misuse. *Canadian Medical Association Journal* 2014; 186(9):341-341.
15. Zhang L, Zheng A, Nie H, Bhavsar K, Xu Y, Sliney DH, Trokel SL, Tsang SH. Laser-induced photic Injury phenocopies macular dystrophy. *Ophthalmic Genetics* 2016; 37:59-67.
16. Combillet F, Saunier V, Rougier MB, Delyfer MN, Korobelnik JF. Multimodal imaging in a case of self-inflicted laser-induced maculopathy. *European Journal Ophthalmology* 2016 26(6): e155 - e157
17. Dhrami-Gavazi E, Lee W, Balaratnasingam C, Kayserman L, Yannuzzi LA, Freund KB. Multimodal imaging documentation of rapid evolution of retinal changes in handheld laser-induced maculopathy. *International Journal of Retina and Vitreous* 2015;1: 14.

## Retinal burns from laser pointers

18. Dolz-Marco R, Cunha Souza E, Lida T, Moreira CA Jr, Nakashima A, Hasegawa T, Freund KB. Iris Atrophy: A Novel Sign of Repeated Self-inflicted Laser Pointer Maculopathy. *Retina* 2016 Oct 26. [Epub ahead of print] Available at [http://journals.lww.com/retinajournal/Citation/publishahead/Iris\\_Atrophy\\_A\\_Novel\\_Sign\\_of\\_Repeated.97237.aspx](http://journals.lww.com/retinajournal/Citation/publishahead/Iris_Atrophy_A_Novel_Sign_of_Repeated.97237.aspx) (Accessed: 05/06/2018)
19. Gosling DB, O'Hagan JB, Quhill FM. Blue Laser Induced Retinal Injury in a commercial pilot at 1300ft. *Aerospace Medicine and Human Performance* 2016; 87:69-70.
20. Hossein M, Bonyadi J, Soheilian R, Soheilian M, Peyman GA. SD-OCT Features of Laser Pointer Maculopathy Before and After Systemic Corticosteroid Therapy. *Ophthalmic Surgery, Lasers & Imaging* 2011; 42:135-138.
21. Joubert R, Farguette F, Chevreaud O, Chiambaretta F, Souied EH. Unilateral maculopathy in a young male patient: A photic laser injury. *Journal français d'ophtalmologie* 2016; 39:786-790.
22. Israeli D, Hod Y, Geyer O. Retinal injury induced by laser pointers. *Harefuah* 2001;140:28-29.
23. Lally DR, Duker JS. Foveal injury from a red laser pointer. *JAMA Ophthalmology* 2014; 132(3):297.
24. Lim ME, Suelzer J, Moorthy RS, Vemuri G. Thermal macular injury from a 154 mW green laser pointer. *Journal of AAPOS* 2014; 18(6):612-614.
25. Luttrull JK, Hallisey J. Laser Pointer-Induced Macular Injury. *American Journal Ophthalmology* 1999;127:95-96.
26. Marshall J, O'Hagan JB, Tyrer JR. Eye hazards of laser 'pointers' in perspective *Br J Ophthalmol* 2016;100:5 583-584

## Retinal burns from laser pointers

27. Palakkamanil MM, Fielden MP. Effects of malicious ocular laser exposure in commercial airline pilots. *Canadian Journal of Ophthalmology* 2015; 50:429-432.
28. Petrou P, Patwary S, Banerjee PJ, Kirkby GR. Bilateral macular hole from a handheld laser pointer. *The Lancet* 2014; 383(9930):1780-1780.
29. Pollithy S, Ach T, Schaal KB, Dithmar S. Acute bilateral impaired vision with central scotoma in an 11-year-old boy. *Ophthalmology* 2012; 109(9):907-910.
30. Raoof N, O'Hagan J, Pawlowska N, Quhill FM. 'Toy' laser macular burns in children: 12-month update. *Eye* 2016; 30:492-496.
31. Raoof N, Bradley P, Theodorou M, Moore AT, Michaelides M. The New Pretender: A Large UK Case Series of Retinal Injuries in Children Secondary to Hand-Held Lasers. *American Journal of Ophthalmology* 2016; 171:88-94.
32. Rusu I, Sherman J, Gallego-Pinazo R, Lam M, Freund KB. Spectral-domain optical coherence tomography and fundus autofluorescence findings in a case of laser pointer-induced maculopathy. *Retinal Cases & Brief Reports* 2013; 7(4):371-375.
33. Sánchez-Barahona, González-Martín-Moro J, Zarallo-Gallardo J, Lozano Escobar I, Cobo-Soriano R. Early changes in optic coherence tomography in a child with laser pointer maculopathy. *Archives de la sociedad Española de Oftalmología* 2016;92: 33-36.
34. Sell CH, Bryan JS. Maculopathy from handheld diode laser pointer. *Archives Ophthalmology* 1999; 117(11):1557-1558.
35. Sethi CS, Grey RH, Hart CD. Laser pointers revisited: a survey of 14 patients attending casualty at the Bristol Eye Hospital. *British Journal Ophthalmology* 1999; 83(10):1164-1167.

## **Retinal burns from laser pointers**

36. Shenoy R, Bialasiewicz AA, Bandara A, Isaac R. Retinal Damage from Laser Pointer Misuse - Case Series from the Military Sector in Oman. *Middle East African Journal Ophthalmology* 2015; 22(3):399-403.
37. Sheyman AT, Nesper PL, Fawzi AA, Jampol LM. Adaptive Optics Imaging in Laser Pointer Maculopathy. *Ophthalmic Surgery, Lasers & Imaging Retina* 2016; 47:782-785.
38. Simonett JM, Scarinci F, Labriola LT, Jampol LM, Goldstein DA, Fawzi AA. A case of recurrent, self-inflicted handheld laser retinopathy. *Journal of AAPOS* 2016;20:168-170.
39. Thanos S, Bohm MRR, Zu HMM, Schmidt PF. Retinal damage induced by mirror-reflected light from a laser pointer. *BMJ Case Reports* 2015, doi:10.1136/bcr-2015-210311.
40. Ueda T, Kurihara I, Koide R. A case of retinal light damage by green laser pointer (Class 3b). *Japanese Journal Ophthalmology* 2011; 55: 428-430.
41. Weng CY, Baomal CR, Albini TA, Berrocal AM. Self-Induced Laser Maculopathy in an Adolescent Boy Utilizing a Mirror. *Ophthalmic Surgery, Lasers & Imaging Retina* 2015; 46:485-488.
42. Wong R, Sim D, Rajendram R, Menon G. Class 3A laser pointer-induced retinal damage captured on optical coherence tomography. *Acta Ophthalmologica Scandinavica* 2007; 85(2):227-228.
43. Wyrsh S, Baenninger PB, Schmid MK. Retinal injuries from a handheld laser pointer. *New England Journal Medicine* 2010; 363(11):1089-1091.

## Retinal burns from laser pointers

44. Xu K, Chin EK, Quiram PA, Davies JB, Parke III DW and Almeida DRP. Retinal Injury Secondary to Laser Pointers in Pediatric Patients. *Pediatrics* Oct 2016, 138 (4) e20161188; DOI: 10.1542/peds.2016-1188 Available at: <http://pediatrics.aappublications.org/content/138/4/e20161188.long> [Accessed: 05/06/2018)
45. Zamir E, Kaiserman I, Chowers I. Laser Pointer Maculopathy. *American Journal of Ophthalmology* 1999; 127:728-729.
46. Ziahosseini K, Doris JP, Turner GS. Laser eye injuries. Maculopathy from handheld green diode laser pointer. *BMJ* 2010; 340:c2982.
47. Quhill F, Quhill H, Pawlowska N. Macular Injury due to Laser Devices: The Results of a Survey Among Consultant Ophthalmologists in the United Kingdom. PO423 - 2016. Presented on Monday October 17, 2016 12:30PM. American Academy of Ophthalmology Congress, Chicago.
48. British standard BS EN 60825-1:2014. *Safety of laser products. Equipment classification and requirements*. British Standards Institution. 2014  
<https://www.thenbs.com/PublicationIndex/documents/details?Pub=BSI&DocID=320928> (Accessed: 05/06/2018)
49. Health Risks from the Use of Laser Pointers. *WHO Information Fact Sheets. Fact Sheet No 202, July 1998* <http://www.who.int/uv/resources/fact/fs202laserpointers.pdf> (Accessed: 05/06/2018)
50. Laser Radiation: Safety Advice. Public Health England August 2014. Available online at: <https://www.gov.uk/government/publications/laser-radiation-safety-advice/laser-radiation-safety-advice> (Accessed: 05/06/2018)



## Retinal burns from laser pointers

51. Performance Standards for Light-Emitting Products, U.S. Food and Drug Administration Code of Federal Regulations, Title 21, Chapter I, Subchapter J, Part 1040, 2016. <http://www.ecfr.gov/cgi-bin/text-idx?SID=c656f500beebfa4cf6a442a1f8f13677&mc=true&node=pt21.8.1040&rqn=div5>.

U.S. Government Publishing Office Bookstore Washington **DC**. (Accessed: 05/06/2018)

52. Important Information for Laser Pointer Manufacturers. *U.S. Food and Drug Administration* 2015 <http://www.fda.gov/Radiation-EmittingProducts/RadiationEmittingProductsandProcedures/HomeBusinessandEntertainment/LaserProductsandInstruments/ucm116373.htm> (Accessed: 05/06/2018).

53. Lee MH, Fox K, Goldwasser S, Lau DWM, Aliahmad B, Sarossy M. Green lasers are beyond power limits mandated by safety standards. Proceedings of the 38th Annual International Conference of the Institute of Electrical and Electronics Engineers (IEEE) Engineering in Medicine and Biology Society (EMBC) 2016 Available at <http://ieeexplore.ieee.org/document/7591885/> (Accessed: 05/06/2018).

54. Hadler J. and Dowell M. Accurate, inexpensive testing of laser pointer power for safe operation. *Measurement Science and Technology* 2013; 24(4):045202.

55. Swift C, Clarkson DE, Allan D, Macpherson A. An assessment of the hazards of 'blue' laser pointers. *Proceedings of 14<sup>th</sup> International Congress of International Radiation Protection Association*. 2016

56. Newson E, Le Maréchal K, David C. Pathological Demand Avoidance Syndrome: a necessary distinction within the pervasive developmental disorders. *Archives of Disease in Childhood* 2003; 88(7):595-600.

57. O'Nions E, Viding E, Greven CU, Ronald, A, Happé F. Pathological demand avoidance: Exploring the behavioural profile. *Autism* 2014; 18(5):538-544.

## Retinal burns from laser pointers

58. ADHD: Clinical Practice Guideline for the Diagnosis, Evaluation, and Treatment of Attention-Deficit/Hyperactivity Disorder in Children and Adolescents. *American Academy of Paediatrics* 2011; 128(5):1007-1022.
59. Minshawi NF, Hurwitz S, Morriss D, McDougale CJ. Multidisciplinary Assessment and Treatment of Self-Injurious Behavior in Autism Spectrum Disorder and Intellectual Disability: Integration of Psychological and Biological Theory and Approach. *Journal of Autism and Developmental Disorders* 2015; 45(6):1541-1568.
60. Weiss JA. Self-injurious behaviours in autism: A literature review. *Journal on Developmental Disabilities* 2002; 9:129-144.
61. Richards C, Moss J, Nelson L, Oliver C. Persistence of self-injurious behaviour in autism spectrum disorder over 3 years: a prospective cohort study of risk markers. *Journal of Neurodevelopmental Disorders*. 2016;8:21. doi:10.1186/s11689-016-9153-x
62. Patton N. Self-inflicted eye injuries: a review. *Eye* 2004;18:867-872.
63. Lee Y, Lenhart PD, Lambert SR. Cataract secondary to self-inflicted blunt trauma in children with autism spectrum disorder. *Journal of AAPOS* (2016), doi: 10.1016/j.jaapos.2016.02.13
64. Bartsch DU, Muftuoglu IK, Freeman WR. Laser Pointers Revisited. *Retina* 2016;36(9):1611-3.
65. Thaung J, Lopes C, Löfgren S. Retinal injuries from handheld lasers: An updated report. Swedish Radiation Safety Authority. Report number: 2015:54 ISSN: 2000-0456 <https://www.stralsakerhetsmyndigheten.se/contentassets/a945e4f1e4c84688b929dedf29fa79dc/201554-retinal-injuries-from-handheld-lasers-an-updated-report> (Accessed: 05/06/2018)

## Retinal burns from laser pointers

66. Public Health England (2016) *The dangers of lasers*. Available at:

<https://www.youtube.com/watch?v=mOs1i4Wxdh4&feature=youtu.be> (Accessed: 05/06/2018).

67. Suffolk Trading Standards. Laser Pointer Safety Tips. July 2016. Available at

<https://suffolktradingstandards.wordpress.com/2016/07/25/laser-pointer-safety-tips/> (Accessed: 05/06/2018).

68. UK Parliament. Laser Misuse (Vehicles) Act 2018. Available at:

<http://www.legislation.gov.uk/ukpga/2018/9/contents/enacted/data.htm> (Accessed: 05/06/18)

69. Department for Business, Energy & Industrial Strategy. Call For Evidence: Laser Pointers. Government Response. Department for Business, Energy & Industrial Strategy, London. January 2018. Available at:

[https://www.gov.uk/government/uploads/system/uploads/attachment\\_data/file/672013/laser-pointers-call-for-evidence-government-response.pdf](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/672013/laser-pointers-call-for-evidence-government-response.pdf) (Accessed: 09/06/2018)

70. Hookham M and Earl B. Seller convicted after laser harms boy's eyesight *The Sunday Times* July 17th 2016 Available <http://www.thetimes.co.uk/article/seller-convicted-after-laser-harms-boys-eyesight-68k08pjlh> (Accessed: 09/06/2018).